

夏の北極振動の異常発達とブロッキング高気圧

立花義裕^{1,2}, 中村哲³

¹ 海洋研究開発機構

² 三重大学大学院生物資源学研究所

³ 国立環境研究所

Abrupt evolution of the summer Arctic Oscillation and its association with blocking

Yoshihiro Tachibana^{1,2} and Tetsu Nakamura³

¹Japan Agency for Marine-Earth Science and Technology, Yokosuka

²Climate and Ecosystem Dynamics Division, Fac. of Bioresources, Mie University

³Atmospheric Environment Division, National Institute for Environmental Studies

We used NCEP-NCAR reanalysis data for the period from 1958 to 2005 to statistically analyze the relationships of the summer Arctic Oscillation (summer AO) with hemispheric scale anomalous summer weather and the occurrence of blocking highs. The anomalous positive AO pattern (low pressure anomaly in the Arctic and high pressure anomaly in midlatitudes) accounts well for the hemispheric scale weather associated with anomalous blocking between the polar and subtropical jets, whereas blocking rarely occurs during negative AO periods. The double jet stream structure is more evident during periods of anomalous positive AO than during periods of negative AO. The surface temperatures associated with the anomalous positive AO clearly show Europe to be hot and Japan to be cool, as was the case during the anomalous summer of 2003. The occurrence of a positive summer AO is therefore consistent with the hemispheric scale anomalous summer weather associated with blocking in 2003. We investigated the abrupt evolution of atmospheric patterns and the geographic distribution of blocking highs associated with the development, maintenance, and decay periods of an anomalous positive AO. During the development period, blocking tends to occur over Europe and the Atlantic Ocean, but no significant blocking signature is evident over eastern Eurasia. During the maintenance stage, blocking tends to occur in the Far East. During the decay stage, blocking over the Pacific region is obvious. This longitudinal migration of blocking phenomena may be used to predict the evolution through time of the NAM. This study is based on Tachibana et al., (2010) published in *the Journal of Geophysical Research*.

References

Tachibana, Y., T. Nakamura, H. Komiya, and M. Takahashi (2010), Abrupt evolution of the summer Northern Hemisphere annular mode and its association with blocking, *J. Geophys. Res.*, 115, D12125, doi:10.1029/2009JD012894.