

スバルバルのコケツンドラに生育するコケ 5 種の光合成特性

内田雅己¹、神田啓史¹、増本翔太²、中坪孝之³

¹ 国立極地研究所

² 横浜国立大学

³ 広島大学

Photosynthetic characteristics of five moss species in a moss tundra on Svalbard

Masaki Uchida¹, Hiroshi Kanda¹, Shota Masumoto² and Takayuki Nakatsubo³

¹National Institute of Polar Research

²Yokohama National University

³Hiroshima University

Arctic terrestrial ecosystems are vulnerable to climate change, and a major concern is how the carbon balance of the ecosystems will respond to climate change (Schuur et al., 2015). Wet land of tundra ecosystems have been strong sinks for atmospheric CO₂ after last glacial and contain large amounts of accumulated soil organic carbon (SOC)(Lund et al., 2010). Climate change is likely to have huge influence on this SOC pool by changing carbon balance. However little known about carbon cycle in the High Arctic wet land ecosystem. As a part of a carbon cycle study in a wet tundra (moss tundra), we investigated relationship between photosynthesis and abiotic factors for five moss species.

We set a study site at Stuphallet where on the Brøgger peninsula near Ny-Ålesund, Svalbard, Norway (79°N). The area is almost totally covered with mosses including relatively dominant species *Calliergon richardsonii*, *Campylopus sp.*, *Campyliadelphus stellatus*, *Sanionia uncinata*, and *Tomenthypnum nitens*. A few vascular plants grow in mosses. We collected a green part of five moss species (*Calliergon richardsonii*, *Meesia triquetra*, *Sanionia uncinata*, *Tomenthypnum nitens*, and *Paludella squarrosa*) and brought back to the NIPR station in Ny-Ålesund. The rates of net photosynthesis and dark respiration for the mosses were determined using an open-flow gas exchange system with an infrared gas analyzer. During the measurements, temperature (2–23°C) and Photosynthetically Active Radiation (0–1500 μmol photons m⁻² s⁻¹) were changed. After the measurements, the moss samples were dried out to determine dry weight. In order to know environmental condition, temperature of green part of moss layer and PAR were also measured at Stuphallet and on the roof of the station, respectively.

During the growing season in 2014 (9 July – 31 Aug), temperature for green part of moss layer ranged from 0 to 16°C. The highest frequency in the temperature was observed at 7°C. 76% of the temperature frequency ranged from 2 to 8°C. On the other hand, Frequency of PAR showed the highest from 100 to 200 μmol photons m⁻² s⁻¹ (26% of the total). The frequency decreased exponentially with increasing PAR more than 200 μmol photons m⁻² s⁻¹ to 1300 μmol photons m⁻² s⁻¹.

Net photosynthetic rate for all moss species within the range of 2 to 12°C did not change significantly when PAR was more than 200 μmol photons m⁻² s⁻¹. In contrast, the photosynthesis tended to decrease with increasing temperature under 100 μmol photons m⁻² s⁻¹. On the other hand, light (PAR) dependence on net photosynthesis tended to be different among the five species. Light saturation point of *C. richardsonii*, *M. triquetra*, and *P. squarrosa* showed approximately 400 μmol photons m⁻² s⁻¹. In contrast, the point for *S. uncinata* and *T. nitens* showed approximately 200 μmol photons m⁻² s⁻¹. In comparison to the net photosynthesis per unit moss dry weight between the five moss species at the most likely temperature and PAR condition (7°C, 200 μmol photons m⁻² s⁻¹), the highest value showed *P. squarrosa* and the lowest showed *T. nitens*. There was more than twice the difference between them. On the other hand, *S. uncinata* and *M. triquetra* showed the highest net photosynthesis at the same temperature and PAR condition per unit area basis. Although net photosynthesis of *P. squarrosa* per unit dry weight basis showed the highest value, the moss per unit area basis showed the lowest value.

The net photosynthesis per unit area at 7°C and 200 μmol photons m⁻² s⁻¹ did not completely coincide with degree dominancy of vegetation at the study area. However, higher degree dominancy of mosses tended to be higher net photosynthesis per unit area. As a next step, we will measure respiration rates for brown part of these mosses and estimate net primary production.