

Tracing meteorite terrestrial weathering effects on rare earth elements (REEs) distribution: implications for meteorite weathering related to climate

Hamed Pourkhorsandi^{1,2}, Vinciane Debaille¹, Massimo D’Orazio², Pierre Rochette³, Rosalind Armytage⁴, and Jérôme Gattacceca³

¹*Laboratoire G-TIME, Université libre de Bruxelles, CP 160/02, 50, Av. F.D. Roosevelt, 1050 Brussels, Belgium*

²*AMU/CNRS, CEREGE, Aix-en-Provence, France*

³*Dipartimento di Scienze della Terra, Università di Pisa, Pisa Italy*

⁴*Jacobs-NASA, Johnson Space Center, Houston, Texas 77058, USA*

Following their fall to Earth, meteorites experience weathering. During this process, their mineralogy and chemical composition can be deeply modified (e.g., Bland et al., 2006). In this study, we focus on REE modifications, owing to their importance in studying meteorite petrogenesis and provenance. Our goal is to observe the effects of weathering on REE composition of the most common meteorite types, the ordinary chondrites (OCs). Meteorites from Atacama (Chile) and Lut (Iran) deserts were selected. Microscopic observations to study the alteration mineralogy were conducted. Then, the REE concentrations and Nd isotope compositions of the samples were measured using ICP-MS and MC-ICP-MS.

Our data show the modification of REE distribution during weathering (Pourkhorsandi et al., 2017). LREE/HREE ratio is typically higher than one and most of the desert meteorites show elevated Σ REE contents with respect to fresh OCs. Despite being heavily weathered at the microscopic scale, meteorites from Lut have less modified REE compositions than those of the moderately weathered meteorites from Atacama. Higher REE modification in the Atacama OCs than those from the Lut desert is interpreted as due to the longer interaction of Atacama meteorites with the terrestrial environment in a drier climate.

Hydrated secondary mineralogy actually prevents the progression of alteration inwards because of the higher volume of secondary minerals, hence acting as a barrier. An important outcome of this study is that even “freshlooking” meteorites can have a chemically altered character (Pourkhorsandi et al., 2017). In addition to hot desert meteorites, we will present data on REE contents of Antarctic OCs. We will also discuss the implications of terrestrial weathering on the Nd isotope systematics in those samples. These samples are being used in cosmochemical studies and a special care should be taken into account while working with them.

References

- Bland, P. A., M. E. Zolensky, G. K. Benedix, and M. Sephton, Weathering of chondritic meteorites. In *Meteorites and the early solar system II*, edited by Lauretta D. and McSween H. Y. Tucson, Arizona: University of Arizona Press., 853–867, 2006.
- Pourkhorsandi, H., M. D’Orazio, P. Rochette, M. Valenzuela, J. Gattacceca, H. Mirnejad, B. Sutter, A. Hutzler, and M. Aboulahris, Modification of REE distribution of ordinary chondrites from Atacama (Chile) and Lut (Iran) hot deserts: Insights into the chemical weathering of meteorites, *Meteoritics and Planetary Science*, 52, 1843-1558.