

Potential diesel and phenol biodegradation by the Antarctic bacterium *Rhodococcus* sp. strain AQ5-07

Siti Aqlima Ahmad^{1,2*}, Nur Adeela Yasid^{1,2}, Gillian Li Yin Lee¹, Ahmad Fareez Ahmad Roslee¹, Nur Nadhirah Zakaria¹, Peter Convey³ and Mohd Yunus Shukor¹.

¹Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

²National Antarctic Research Centre, B303 Level 3, Block B, IPS Building, Universiti Malaya, 50603 Kuala Lumpur, Malaysia

³British Antarctic Survey, NERC, High Cross, Madingley Road, Cambridge CB3 0ET, United Kingdom

The potential study of Antarctic microorganisms in bioremediation has been of increasing interest, due to their adaptations to harsh conditions and their metabolic potential in removing a wide variety of organic pollutants at low temperature. However, there is currently inadequate information available about indigenous Antarctic diesel and phenol degraders. Environmental variables such as temperature, pH and salinity are important factors affecting the efficiency of diesel and phenol degradation (Ruberto et al., 2005; Habib et al., 2018). In this study, a psychrophilic bacterium from King George Island, Antarctica (Figure 1) was isolated and identified as *Rhodococcus* sp. strain AQ5-07 by 16S rRNA analysis (Figure 2) (Lee et al., 2018). The optimum conditions of strain AQ5-07 for diesel and phenol degradation were optimized (Table 1). For diesel degradation, strain AQ5-07 displayed optimal cell growth and biodegradation activity at 1% (v/v) of initial diesel concentration, 15-20°C, pH 7 and 0.1 g/L NaCl. While, it optimal phenol degradation occurred at 0.5 g/L phenol, 10-15°C, pH 7.0 and 0.15 g/L NaCl. Based on the results obtained, strain AQ5-07 show good potential as a candidate for biodegradation operations on diesel and phenol contaminated sites in polar region, especially in Antarctica.




Figure 1. Map of sampling location, King George Island, Antarctica.




Figure 2. Maximum likelihood phylogenetic tree of strain AQ5-07 with 20 related type strains of *Rhodococcus* spp. based on 16S rRNA sequences. *Escherichia coli* was used as an outgroup.

Table 1. Comparison of result from OFAT on diesel and phenol degradation.

Parameter	Optimization	
	Diesel degradation (1%)	Phenol degradation (0.5 g/L)
Temperature (°C)	15-20	10-15
pH	7	7
Salinity (NaCl concentration mg/L)	0.1	0.15

References

Habib, S., Ahmad, S.A., Wan Johari, W.L., Shukor, M.Y., Alias, S.A., Khalil, K.A. and N.A. Yasid, Evaluation of conventional and response surface level optimisation of n-dodecane (n-C12) mineralisation by psychrotolerant strains isolated from pristine soil at Southern Victoria Island, Antarctica. *Microbial Cell Factories*, 17 (44), doi 10.1186/s12934-018-0889-8, 2018.

Lee, G.L.Y., Ahmad, S.A., Yasid, N.A., Zulharnain, A., Convey, P., Wan Johari, W.L., Alias, S.A, Gonzalez-Rocha, G. and M.Y. Shukor, Biodegradation of phenol by cold-adapted bacteria from Antarctic soils. *Polar Biology*, 41, 553-562, 2018.

Ruberto, L.A.M., Vazquez, S., Lobalbo, A. and W.P. MacCormack, Psychrotolerant hydrocarbon-degrading *Rhodococcus* strains isolated from polluted Antarctic soils. *Antarctic Science* 17, 47-56, 2005.