

# Petrology of orthogneisses from the Mozambique Belt in southern Malawi: Preliminary report

Toshiaki Tsunogae<sup>1,2</sup>, Sam Uthup<sup>1</sup>, Mzee Wandembo Nyirongo<sup>3</sup>, Yusuke Takamura<sup>1</sup>, Kazuki Takahashi<sup>1</sup> and Md. Sazzadur Rahman<sup>1</sup>

<sup>1</sup>*Graduate School of Life and Environmental Sciences, University of Tsukuba, Japan*

<sup>2</sup>*Department of Geology, University of Johannesburg, South Africa*

<sup>3</sup>*Spring Stone LTD, Zomba, Malawi*

Detailed petrological and geochronological studies on Pan-African high-grade metamorphic rocks in East Antarctica, India, Sri Lanka, Madagascar, and East Africa for the last several years confirmed that the regions were formed by complex subduction and collisional processes related to the amalgamation of Gondwana Supercontinent during latest Neoproterozoic to Cambrian (e.g., Santosh et al., 2017; Kazami et al., 2016; Takamura et al., 2016, 2018; He et al., 2016; Takahashi et al., 2017, 2018; Tsunogae et al., 2015, 2016; Kuribara et al., 2018; Tang et al., 2018). The basement rocks in southern Malawi form part of the Mozambique Belt (or the southern Irumide Belt) which correspond a typical example of Neoproterozoic collisional orogens in southeastern Africa (e.g., Kröner et al., 2001). Southern Malawi is particularly important because the region corresponds to the junction between the East-African Orogen (ca. 600-550 Ma) and the Kuunga orogen (ca. 560-530 Ma), therefore complex magmatic/metamorphic episodes might have been recorded in high-grade metamorphic rocks exposed in this region. Below we summarize the results of our preliminary geological, petrological, and geochemical studies based on our fieldwork in southern Malawi in July 2018.

The dominant lithologies in Lilongwe-Zomba-Blantyre area in southern Malawi are biotite gneiss (biotite + quartz + microcline + plagioclase), charnockite (orthopyroxene + K-feldspar + quartz + plagioclase), mafic granulite (plagioclase + clinopyroxene + orthopyroxene + garnet + hornblende + quartz), and quartzite which are intruded by syenite (K-feldspar + biotite + calcic amphibole + quartz) and granite (quartz + K-feldspar + plagioclase + biotite). Major, minor, and REE geochemical data for selected felsic orthogneisses suggest arc-magmatic signatures of the rocks, whereas mafic orthogneisses shows both magmatic arc and non-arc signatures. Granulite-facies conditions of 750-810 °C were obtained based on conventional geothermometry of garnet-bearing mafic granulites. The results of this study suggest that the southern part of Malawi with dominant orthogneisses corresponds to a Neoproterozoic (1.0 Ga; Kröner et al., 2001) magmatic arc which was metamorphosed during the latest Neoproterozoic collision of the Kalahari and the Congo Cratons.

## Acknowledgement

This study was partly supported by a Grant-in-Aid for Scientific Research (B) from Japan Society for the Promotion of Science (JSPS) (No. 18H01300) to Tsunogae.

## References

- He, X.-F., M. Santosh, T. Tsunogae, S.P.K. Malaviarachchi and P.L. Dharmapriya, Neoproterozoic arc accretion along the 'eastern suture' in Sri Lanka during Gondwana assembly, *Precambrian Research*, 279, 57-80, 2016.
- Kazami, S., T. Tsunogae, M. Santosh, Y. Tsutsumi and Y. Takamura, Petrology, geochemistry and zircon U-Pb geochronology of a layered igneous complex from Akarui Point in the Lützow-Holm Complex, East Antarctica: Implications for Antarctica-Sri Lanka correlation, *Journal of Asian Earth Sciences*, 130, 206-222, 2016.
- Kröner, A., A.P. Willner, E. Hegner, P. Jaekel and A. Nemchin, Single zircon ages, PT evolution and Nd isotopic systematics of high-grade gneisses in southern Malawi and their bearing on the evolution of the Mozambique belt in southeastern Africa, *Precambrian Research*, 109, 257-291, 2001
- Kuribara, Y., T. Tsunogae, Y. Takamura and Y. Tsutsumi, Petrology, geochemistry, and zircon U-Pb geochronology of the Zambezi Belt in Zimbabwe: Implications for terrane assembly in southern Africa, *Geoscience Frontiers*, doi: 10.1016/j.gsf.2018.05.019, 2018.
- Santosh, M., C.-N. Hu, X.-F. He, S.-S. Li, T. Tsunogae, E. Shaji and G. Indu, Neoproterozoic arc magmatism in the southern Madurai Block, India: Subduction, reamination, continental outbuilding, and the growth of Gondwana, *Gondwana Research*, 45, 1-42, 2017.
- Takahashi, K. and T. Tsunogae, Carbonic fluid inclusions in a garnet-pyroxene granulite from Austhovde in the Lützow-Holm Complex, East Antarctica: Implications for a decompressional *P-T* path, *Journal of Mineralogical and Petrological Sciences*, 112, 132-137, 2017.

- Takahashi, K., T. Tsunogae, M. Santosh, Y. Takamura and Y. Tsutsumi, Paleoproterozoic (ca. 1.8 Ga) arc magmatism in the Lützow-Holm Complex, East Antarctica: implications for crustal growth and terrane assembly in erstwhile Gondwana fragments, *Journal of Asian Earth Sciences*, 157, 245-268, 2018.
- Takamura, Y., T. Tsunogae, M. Santosh, S.P.K. Malaviarachchi and Y. Tsutsumi, U-Pb geochronology of detrital zircon in metasediments from Sri Lanka: Implications for the regional correlation of Gondwana fragments, *Precambrian Research*, 281, 434-452, 2016.
- Takamura, Y., T. Tsunogae, M. Santosh and Y. Tsutsumi, Detrital zircon geochronology of the Lützow-Holm Complex, East Antarctica: Implications for Antarctica - Sri Lanka correlation, *Geoscience Frontiers*, 9, 355-375, 2018.
- Tang, L., M. Pan, T. Tsunogae, Y. Takamura and Y. Tsutsumi, Y, Cordierite-bearing granulites from Ihosy, southern Madagascar: Petrology, geochronology and regional correlation of suture zones in Madagascar and India, *Geoscience Frontiers*, doi: 10.1016/j.gsf.2018.05.014, 2018.
- Tsunogae, T., Q.Y. Yang and M. Santosh, Early Neoproterozoic arc magmatism in the Lützow-Holm Complex, East Antarctica: Petrology, geochemistry, zircon U-Pb geochronology and Lu-Hf isotopes and tectonic implications, *Precambrian Research*, 266, 467-489, 2015.
- Tsunogae, T., Q.Y. Yang and M. Santosh, Neoproterozoic - Early Paleoproterozoic and Early Neoproterozoic arc magmatism in the Lützow-Holm Complex, East Antarctica: insights from petrology, geochemistry, zircon U-Pb geochronology and Lu-Hf isotopes, *Lithos*, 263, 239-256, 2016.