

## Black Carbon in Snowpack over the Different Regions in the Arctic

Tatsuhiko Mori<sup>1</sup>, Kumiko Goto-Azuma<sup>2,3</sup>, Yutaka Kondo<sup>2</sup>, Yoshimi Ogawa-Tsukagawa<sup>2</sup>, Kazuhiko Miura<sup>1</sup>, Motohiro Hirabayashi<sup>2</sup>, Makoto Koike<sup>4</sup>, Nobuhiro Moteki<sup>4</sup>, Sho Ohata<sup>4</sup>, Naga Oshima<sup>5</sup>, P. R. Sinha<sup>6</sup>, Konosuke Sugiura<sup>7</sup>, Teruo Aoki<sup>8</sup>, Martin Schneebeli<sup>9</sup>, Konrad Steffen<sup>9</sup>, Atsushi Sato<sup>10</sup>, Akane Tsushima<sup>11</sup>, Vladimir Makarov<sup>12</sup>, Satoshi Omiya<sup>13,14</sup>, Atsuko Sugimoto<sup>15</sup>, and Shinya Takano<sup>16</sup>

<sup>1</sup>*Tokyo University of Science*, <sup>2</sup>*National Institute of Polar Research*, <sup>3</sup>*The Graduate University for Advanced Studies*, <sup>4</sup>*Graduate School of Science, The University of Tokyo*, <sup>5</sup>*Meteorological Research Institute*, <sup>6</sup>*Balloon Facility, Tata Institute of Fundamental Research*, <sup>7</sup>*University of Toyama*, <sup>8</sup>*Department of Earth Sciences, Faculty of Science, Okayama University*, <sup>9</sup>*WSL Institute for Snow and Avalanche Research*, <sup>10</sup>*former Snow and Ice Research Center*, <sup>11</sup>*Reserach Institute for Humanity and Nature*, <sup>12</sup>*Melnikov's Permafrost Institute*, <sup>13</sup>*Institute of Low Temperature Science, Hokkaido University*, <sup>14</sup>*Civil Engineering Research Institute for Cold Region*, <sup>15</sup>*Arctic Research Center, Hokkaido University*, <sup>16</sup>*Graduate School of Environmental Science, Hokkaido University*.

Black carbon (BC) deposited on snow lowers snow albedo, potentially contributing to the warming in the Arctic. It is critically important to measure the spatial distributions of BC in snowpack in different regions of the Arctic to quantify this effect. However, accurate measurements of BC in snowpack are very limited because of the large uncertainties in the previous measurements. We measured size-resolved BC mass concentrations ( $C_{\text{MBC}}$ ) in snowpack over Finland, Alaska, Siberia, Greenland, and Spitsbergen in early spring between 2012 and 2016 by using a single-particle soot photometer. We collected 108 samples of surface snow, 77 samples of subsurface snow, and 112 columns of snowpack in these regions in total. The amounts of BC deposition ( $\text{DEP}_{\text{MBC}}$ ) during snow accumulation periods were derived from  $C_{\text{MBC}}$  and snow water equivalent (SWE). The average  $C_{\text{MBC}}$  in each region was well correlated with the average anthropogenic BC emission flux ( $r^2 = 0.77$ ), suggesting significant influences of anthropogenic BC emissions on  $C_{\text{MBC}}$  on regional scales. The average size distributions of BC in snowpack shifted to smaller sizes with the decrease in  $C_{\text{MBC}}$ , likely due to an increase in the removal efficiency of BC with the increase in BC diameter during the transport from major sources. The previous  $C_{\text{MBC}}$  obtained by using an Integrating Sphere/Integrating Sandwich spectrophotometer were much larger than our  $C_{\text{MBC}}$  in the overlapping regions. The present data of  $C_{\text{MBC}}$ , SWE, and  $\text{DEP}_{\text{MBC}}$  will be useful in constraining climate model to estimate the effect of BC on the climate of the Arctic.