

Concentrations and size distribution of black carbon in Northwest Greenland during the past 350 years reconstructed from an ice core

Kumiko Goto-Azuma^{1,2}, Yoshimi Ogawa-Tsukagawa¹, Yutaka Kondo¹, Remi Dallmayr^{1,*}, Motohiro Hirabayashi¹, Jun Ogata¹, Kyotaro Kitamura¹, Kenji Kawamura^{1,2}, Hideaki Motoyama^{1,2}, Sumito Matoba³, Teruo Aoki⁴, Nobuhiro Moteki⁵, Sho Ohata⁵, Tatsuhiro Mori⁶, Makoto Koike⁵, Yuki Komuro⁷, Akane Tsushima⁸, Naoko Nagatsuka¹, Wataru Shigeyama^{1,2} and Koji Fujita⁸

¹*National Institute of Polar Research, Japan*

²*SOKENDAI (The Graduate University for Advanced Studies), Japan*

³*Hokkaido University, Japan*

⁴*Okayama University, Japan*

⁵*University of Tokyo, Japan*

⁶*Tokyo University of Science, Japan*

⁷*Yamagata University, Japan*

⁸*Nagoya University, Japan*

** Present affiliation: Alfred Wegner Institute for Polar and Marine Research, Germany*

An ice core to the depth of 225 m was drilled at the SIGMA-D site, Northwest Greenland, in 2014 under the SIGMA (Snow Impurity and Glacial Microbe Effects on Abrupt Warming in the Arctic) project (Matoba et al., 2015). We analyzed the core to the depth of 113 m. The top 6 m of the core was cut at a ca 5 cm interval, melted and analyzed for black carbon (BC) together with stable isotopes of water, microparticles and six elements (Na, K, Mg, Ca, Fe, and Al). Between 6 and 113 m depths, we used a Continuous Flow Analysis (CFA) system developed at the National Institute of Polar Research (NIPR). The NIPR CFA system allows high resolution analysis of black carbon (BC) together with stable isotopes of water, microparticles, electric conductivity, and six elements (Na, K, Mg, Ca, Fe, and Al). For BC analysis, we used a recently developed Wide-range SP2 (Single Particle Soot Photometer, Droplet Measurement Technologies). While a normal SP2 detects BC particles in the size range between 70 and 800 nm, the Wide Range SP2 enabled us to detect BC particles in the size range between 70 and 4000 nm (Mori et al., 2016).

Here we report the variability of BC concentration and size distribution over the past 350 years. BC concentrations started to increase in the 1890s, reached its maximum in the 1930's - 1940's, and decreased again since then. The increase is likely due to anthropogenic input. We find anthropogenic changes in size distribution of BC: the mode diameter of mass concentration became larger. We also find anthropogenic changes in seasonality of BC concentration: Annual concentration peak shifted from summer to winter.

References

Matoba, S., H. Motoyama, K. Fujita, T. Yamasaki, M. Minowa, Y. Onuma, Y. Komuro, T. Aoki, S. Yamaguchi, S. Sugiyama and H. Enomoto, Glaciological and meteorological observations at the SIGMA-D site, northwestern Greenland Ice Sheet. *Bull. Glaciol. Res.*, 33, 7-10, DOI:10.5331/bgr.33.7 2015.

Mori, T., N. Moteki, S. Ohata, M. Koike, K. Goto-Azuma, Y. Miyazaki and Y. Kondo, Improved technique for measuring the size distribution of black carbon particles in liquid water, *Aerosol Science & Technology*, 50, 3, 242-254, DOI: 10.1080/02786826.2016.1147644, 2016.