

# Metamorphism of layered firn at Dome Fuji, Antarctica: Evolution of relations between Near-infrared reflectivity and the other textural/chemical properties

Shuji Fujita, 1,2,  
Kumiko Goto-Azuma, 1,2  
Motohiro Hirabayashi, 1  
Akira Hori, 3  
Yoshinori Iizuka, 4  
Yuko Motizuki, 5  
Hideaki Motoyama, 1,2,  
Kazuya Takahashi, 5

1National Institute of Polar Research, Research, Organization of Information and Systems (ROIS)  
10-3 Midori-cho, Tachikawa, Tokyo 190-8518, Japan

2Department of Polar Science, The Graduate University for Advanced Studies (SOKENDAI), 10-3  
Midori-cho, Tachikawa,  
Tokyo 190-8518, Japan

3Kitami Institute of Technology, Kitami, Japan

4Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan

5RIKEN Nishina Center, Wako, Japan

Correspondence: S. Fujita <[sfujita@nipr.ac.jp](mailto:sfujita@nipr.ac.jp)>

Evolution of polar firn was investigated at sites at Dome Fuji, to better understand signals of deep ice cores. Using samples from a 4-m-deep pit and a 122-m-deep core, relations between major textural and chemical properties, such as Near-infrared light reflectivity  $R$ , density  $\rho$ , microwave dielectric anisotropy  $\Delta\epsilon$ , and concentration of major ions, were investigated at a depth range of 0 – 122 m, with high spatial resolutions. At the near-surface depths, we found: (i) Fluctuations of  $R$ ,  $\rho$ , and  $\Delta\epsilon$  are positively correlated; (ii)  $\Delta\epsilon$  ranges 0.03 – 0.07 immediately below the snow surface at ~0.1 m depth; (iii) These properties of  $R$ ,  $\rho$ , and  $\Delta\epsilon$  are not correlated to major ions. With increasing depths during reported phenomena of density crossover, the positive correlation of  $R$  to  $\Delta\epsilon$  persistently remains with a slight decrease. Besides,  $R$  becomes weakly negatively correlated to concentration of  $\text{Na}^+$  which is the sea salt marker. These facts suggest that textural features of the near-surface depths are preserved in both  $R$  and  $\Delta\epsilon$  at a depth range immediately below bubble-close-off, being weakly affected by reported softening of ice by  $\text{Cl}^-$  ions. We therefore suggest that optically layered features in ice cores are directly linked to the metamorphism.