

Walking behavior of the Antarctic tardigrade, *Acutuncus antarcticus*

Kurumi Yoshimura¹, Megumu Tsujimoto²

^{1,2} Faculty of Environment and Information Studies, Keio University

According to Vogel (1988), swimming is a better mean of transportation than walking in terms of energy consumption for aquatic animal. Despite having many similarities in their ecology including habitats and tolerance abilities against extreme environments as the Antarctic terrestrial meiofauna, rotifers and nematodes swim, on the other hand, tardigrades walk for the transportation. We wondered whether there are any advantages for Tardigrada to choose walking instead of swimming as a mean of transportation.

In the present research, we observed the walking behavior of the Antarctic tardigrade, *Acutuncus antarcticus*. *Acutuncus antarcticus* is the tardigrade species endemic to Antarctica and widely distributed around the continent (Velasco-Castrillón et al. 2014). Individual tardigrades of the *A. antarcticus* LSW strain (Tsujimoto et al. 2015, 2020) were placed on flat agar plates and their walking behaviors were recorded using Stereomicroscope (Olympus Research Stereomicroscope System SZX10) and attached USB camera (Raymer's USB digital camera for microscopes WrayCAM-NOA2000). We then collected the data of the following four characteristics, 1) walking speed, 2) type of the walk (classification of how they walk), 3) movement during the directional change, and 4) grounding ratio (percentage of the legs which are attached on the plate) and analyzed. Some of the data we collected here on the Antarctic tardigrade will be compared from the data obtained from the previous study on two other temperate species of tardigrades, *Ramazzottius varieornatus* and *Milnesium* sp., and we will try to find the differences in their walking behavior.

References

- Vogel S. *Life's Devices : The Physical World of Animals and Plants*. Princeton University Press, 1988-12-21.
- Velasco-Castrillón A, Gibson JAE, Stevens MI (2014) A review of current Antarctic limno-terrestrial microfauna. *Polar Biol* 37:1517–1531
- Tsujimoto M, Kagoshima H, Kanda H, Watanabe K, Imura S. Reproductive performance of the Antarctic tardigrades *Acutuncus antarcticus* (Eutardigrada: Hypsibiidae) revived after being frozen for over 30 years and of their offspring. *Zoological Journal of the Linnean Society*, 188(3): 839 – 847, 2020.
- Tsujimoto M, Kagoshima H, Kanda H, Watanabe K, Imura S. Reproductive performance of the Antarctic tardigrades *Acutuncus antarcticus* (Eutardigrada: Hypsibiidae) revived after being frozen for over 30 years and of their offspring. *Zoological Journal of the Linnean Society*, 188(3): 839 – 847, 2020.