

Traces of fluid movement in the active lower crust of collision setting observed in the Sør Rondane Mountains, East Antarctica: A field survey report of JARE61

Tetsuo Kawakami¹, Tatsuro Adachi², Masaoki Uno³, Fumiko Higashino⁴ and Noriyoshi Tsuchiya³

¹Kyoto University, ²Kyushu University, ³Tohoku University, ⁴Okayama University of Science

The Sør Rondane Mountains (SRM), East Antarctica, is considered as a collision zone formed at ~650-600 Ma (Osanai et al. 2013), and traces of geofluids in the lower crust of the collision setting can be widely observed (Higashino et al., 2013; Kawakami et al., 2017; Uno et al., 2017). Aim of field survey by the 61st Japan Antarctic Research Expedition (JARE61) SRM geology team was to constrain the pressure-temperature-deformation-time-fluid (*P-T-D-t-fluid*) paths of the high-grade metamorphic rocks widely from the SRM and understand the role of geofluids in the evolution of the crust.

In several map-scale mylonitic shear zones (Oyayubi-one and Mefjell), Cl-rich selvages mainly composed of biotite and hornblende were confirmed by *in situ* handy XRF analysis, suggesting that these shear zones were possibly the pathways of Cl-bearing fluid. In addition to the large-scale shear zones, mm-cm thick black selvages composed of Cl-bearing biotite or hornblende (\pm garnet) are also the traces of Cl-bearing fluid infiltration postdating gneissosity-forming deformation (Higashino et al., 2019). We confirmed wide distribution of such gneissosity-cutting selvages both in NE- and SW-terranes. In most cases, such selvages fill small-scale shear zones that accompany cm- to decimeter-scale displacements. Such selvages were locally developed in the interboudin partitions developed in a garnet-biotite gneiss, suggesting that the timing of the presence of Cl-bearing fluid included the last stage of gneissosity-forming deformation.

Recently, timing of garnet-forming high-grade metamorphism in the SRM was dated to be >600 Ma and <580 Ma both in the NE- and SW-terranes, based on the evaluation of distribution coefficients between metamorphic zircon and garnet (Higashino et al., 2020). This observation suggests that the main metamorphism recorded in high-grade metamorphic rocks could be different in timing from locality to locality, and detailed *P-T-D-t-fluid* path analysis is required for the comprehensive understanding of the geofluid activities in the active lower crust.

References

- Higashino, F., Kawakami, T., Satish-Kumar, M., Ishikawa, M., Maki, K., Tsuchiya, N., Grantham, G., Hirata, T. 2013, Chlorine-rich fluid or melt activity during granulite facies metamorphism in the Late Proterozoic to Cambrian continental collision zone – an example from the Sør Rondane Mountains, East Antarctica. *Precambrian Research*, 234, 229-246.
- Higashino, F., Kawakami, T., Tsuchiya, N., Satish-Kumar, M., Ishikawa, M., Grantham, G., Sakata, S., Hirata, T., 2019, Brine Infiltration in the Middle to Lower Crust in a Collision Zone: Mass Transfer and Microtexture Development Through Wet Grain-Boundary Diffusion, *Journal of Petrology*, 60 (2), 329-358.
- Higashino, F., Kawakami, T., Sakata, S., Hirata, T., 2020, Multiple timings of garnet-forming high-grade metamorphism in the Sør Rondane Mountains, East Antarctica. Abstract of the 11th Symposium on Polar Science, NIPR.
- Kawakami, T., Higashino, F., Skrzypek, E., Satish-Kumar, M., Grantham, G., Tsuchiya, N., Ishikawa, M., Sakata, S., Hirata, T., 2017, Prograde infiltration of Cl-rich fluid into the granulitic continental crust from a collision zone in East Antarctica (Perlebandet, Sør Rondane Mountains), *Lithos*, 274-275, 73-92.
- Osanai, Y., Nogi, Y., Baba, S., Nakano, N., Adachi, T., Hokada, T., Toyoshima, T., Owada, M., Satish-Kumar, M., Kamei, A., Kitano, I., 2013, Geologic evolution of the Sør Rondane Mountains, East Antarctica: Collision tectonics proposed based on metamorphic processes and magnetic anomalies, *Precambrian Research*, 234, 8-29.
- Uno, M., Okamoto, A., Tsuchiya, N., 2017, Excess water generation during reaction-inducing intrusion of granitic melts into ultramafic rocks at crustal P-T conditions in the Sør Rondane Mountains of East Antarctica, *Lithos*, 284-285, 625-641.