

Adaptation strategy of aerial green alga, *Prasiola crispa* growing in Antarctica

Makiko Kosugi¹, Fumino Maruo², Norio Kurosawa³, Akinori Kawamata⁴, Yasuhiro Kamei⁵, Yasuhiro Kashino⁶, Hiroyuki Koike², Sakae Kudoh⁷ and Satoshi Imura⁷

¹*Astro Biology Center*

²*Chuo University*

³*Soka University*

⁴*Ehime Prefectural Science Museum*

⁵*National Institute for Basic Biology (NIBB)*

⁶*University of Hyogo*

⁷*National Institute of Polar Research (NIPR)*

Lichens, bryophyta, algae and cyanobacteria have a critical role as primary producer in terrestrial ecosystem of Antarctica. They are surviving the severe condition exposed to low temperature, desiccation, strong wind and high light stresses. *Prasiola crispa* is one of the dominant terrestrial green algae in Antarctic region, and often constitutes a large sheet-like colony on the soil surface. We set up a year-round micrometeorological observing system at the habitats of *P. crispa*, *Umbilicaria decussate* (lichen) and *Bryum argenteum* (bryophyta) in the Yukidori Zawa of Lnghovde, and obtained data every 10 min for five years (from Jan. 2013 to Feb. 2018). The observing items were temperature, relative humidity (RH), PAR around organisms and organisms' temperatures. On the other hand, for the analysis of physiological aspect, we studied the dependence of wavelengths from 320nm to 750nm against photodamage in photosystem by using the Okazaki Large Spectrograph (National Institute for Basic Biology, Okazaki, Japan)^a.

During the observation period of five years, the monthly RH and PAR showed high variety in summer season (January to March) but temperature showed similar trend except for January. January 2013 was warmest month and the precipitation was so limited. The monthly average temperatures around organisms were higher than 5°C and maximum temperature of organisms over 25°C, whereas the monthly average and the maximum temperatures in general years were approximately 2 and 17°C, respectively. Water availability and light intensity in summer season are expected to have a big impact to net photosynthetic production. We predicted active periods and photo damage risk in the three photosynthetic organisms by using physiological experimental data. As a result, risk of photodamage in *P. crispa* is highest in January and low light condition such as cloudy day is suitable for photosynthesis. On the other hand, in our recent study, it was revealed *P. crispa* had a unique light-harvesting chlorophyll binding protein (Pc-frLHC) which absorbed far-red light and enable to excitation of photosystem II^b. Pc-frLHC is expressed in the lower side of layer of colony and is contributing to increase net photosynthetic production inside of colony. We mention about adaptation strategy of *P. crispa* growing Antarctica from ecological and physiological aspects.

(Supported by grant 24770030 and 17K19431 from the Japan Society for the Promotion of Science and grant 151376 from the Sumitomo Foundation.)

References

^aKosugi, M., Maruo, F., Inoue, T., Kurosawa, N., Kawamata, A., Koike, H., Kamei, K., Kudoh, S. and Imura, S. A comparative study of wavelength-dependent photoinactivation in photosystem II of drought-tolerant photosynthetic organisms in Antarctica and the potential risks of photoinhibition in the habitat., *Annals of Botany*, 122(7) 1263 – 1278, 2018..

^bKosugi, M., Ozawa, S., Takahashi, Y., Kamei, Y., Itoh, S., Kudoh, S., Kashino, Y. and Koike, H. Red-shifted chlorophyll *a* bands allow uphill energy transfer to photosystem II reaction centers in an aerial green alga, *Prasiola crispa*, harvested in Antarctica., *Biochimica et Biophysica Acta - Bioenergetics* 1861 148139, 2020