

Diatoms define a novel freshwater biogeography of the Antarctic

Elie Verleyen¹, Bart Van de Vijver^{2,3}, Bjorn Tytgat¹, Eveline Pinseel^{1,2}, Dominic A. Hodgson⁴, Kateřina Kopalová^{5,6}, Steven L. Chown⁷, Eric Van Ranst⁸, Satoshi Imura^{9,10}, Sakae Kudoh^{9,10}, Wim Van Nieuwenhuyze¹, Koen Sabbe¹, Wim Vyverman¹

¹*Ghent University, Protistology and Aquatic Ecology, Campus de Sterre, Krijgslaan 281 S8, B-9000 Gent, Belgium.*

²*Meise Botanic Garden, Nieuwelaan 38, 1860 Meise, Belgium.*

³*University of Antwerp, Department. Biology – ECOBE, Universiteitsplein 1, B-2610 Antwerp, Belgium.*

⁴*British Antarctic Survey, Natural Environment Research Council, High Cross Madingley Road, CB3 0ET, Cambridge, UK.*

⁵*Charles University, Department of Ecology, Viničná 7, 128 44 Prague 2, Czech Republic.*

⁶*Phycology Centre, Institute of Botany, Academy of Science CR, Dulekská 135, 37982 Třeboň, Czech Republic.*

⁷*School of Biological Sciences, Monash University, Melbourne, VIC 3800, Australia*

⁸*Ghent University, Department of Geology (WE13), Campus de Sterre, Krijgslaan 281 - S8, B-9000 Gent, Belgium.*

⁹*National Institute of Polar Research, 10-3, Midoricho, Tachikawa, Tokyo 190-8518, Japan.*

¹⁰*The Graduate University for Advanced Studies, SOKENDAI, 10-3, Midoricho, Tachikawa, Tokyo 190-8518, Japan.*

Terrestrial biota in the Antarctic are more globally distinct and highly structured biogeographically than previously believed, but information on biogeographic patterns and endemism in freshwater communities is largely lacking. We studied biogeographic patterns of Antarctic freshwater diatoms based on the analysis of species occurrences in a dataset of 439 lakes spread across the Antarctic realm. Highly distinct diatom floras, both in terms of composition and richness, characterize Continental Antarctica, Maritime Antarctica and the sub-Antarctic islands, with marked biogeographic provincialism in each region. A total of 44% of all species is estimated to be endemic to the Antarctic, and most of them are confined to a single biogeographic region. The level of endemism significantly increases with increasing latitude and geographic isolation. Our results have implications for conservation planning, and suggest that successful dispersal of freshwater diatoms to and within the Antarctic is limited, fostering the evolution of highly endemic diatom floras.